

## Original Article

### Comparison of Supraglottic Devices I-gel and LMA Fastrach<sup>R</sup> as Conduit for Endotracheal Intubation

Himanshu Sagar<sup>1</sup>, Gurdeep Singh Jheetay<sup>2</sup>, Ankita Mago<sup>3</sup>, Sukhvinder Singh Oberoi<sup>4</sup>, Raj Kumar<sup>5</sup>

<sup>1</sup>Post-graduate student, <sup>2</sup>Professor and Head, Department of Anaesthesia, <sup>3</sup>Senior Resident, Department of Obstetrics and Gynaecology, Teerthanker Mahaveer Medical College and Research Centre, <sup>4</sup>Professor and Head, Department of Public Health Dentistry, <sup>5</sup>Reader, Department of Microbiology, Genesis Institute of Dental Sciences and Research, Ferozpur, Punjab, India

#### ABSTRACT

**Aim and objectives:** Comparative evaluation of clinical performance of I-GEL with LMA fastrach in elective surgeries. **Materials and method:** The present study was conducted in the Department of Anesthesiology, Teerthanker Mahaveer Medical College & Research Centre, Moradabad during the period 2015-2017, on 80 patients undergoing elective surgery under general anesthesia. The parameters recorded were Time taken for successful placement, Time taken for insertion of Endo-tracheal Tube, Number of insertion attempts, Quality of ventilation during anesthesia, Haemodynamic parameters such as pulse rate, systolic and diastolic blood pressure and SpO<sub>2</sub>, Airway trauma and Gastric distension. **Results:** There was no significant difference in mean Heart Rate, Systolic and Diastolic blood pressure and mean Arterial pressure at Baseline, 1 minute, 3 minutes and 5 minutes between I-GEL and LMA FOSTRACH groups. The mean Time taken for supra-glottic device was significantly more among LMA FOSTRACH (31.57±3.08) in comparison to I-gel (15.07±1.65). The mean Time taken for Endo-tracheal tube was significantly more among I-gel (24.00±0.31) minutes in comparison to LMA FOSTRACH (19.95±0.37). **Conclusion:** Both Fastrach LMA and I-gel are suitable devices to be used as conduit to endotracheal intubation particularly in susceptible patients where hemodynamic disturbances during intubation are not required. But I-gel proved to be more better than Fastrach LMA in terms of benefits.

**Key words:** Elective surgeries, Endo-tracheal Tube, General anesthesia, Hemodynamic disturbances.

Received: 20 January 2018

Revised: 22 February 2018

Accepted: 26 February 2018

**Corresponding author:** Dr. Himanshu Sagar, Post-graduate student, Department of Anaesthesia, Teerthanker Mahaveer Medical College and Research Centre, India

**This article may be cited as:** Sagar H, Jheetay GS, Mago A, Oberoi SS, Kumar R. Comparison of Supraglottic Devices I-gel and LMA Fastrach<sup>R</sup> as Conduit for Endotracheal Intubation. Int J Res Health Allied Sci 2018; 4(2):85-92.

#### INTRODUCTION

Secure airway management in anesthesia is critical for appropriate treatment of respiratory complications and successful operation.<sup>[1]</sup> Successful transition of these devices through anatomical elements and their proper placement is vital.<sup>[2]</sup> Intubation is one of the most important procedures related to prognosis in severely ill patients.<sup>[3]</sup> Tracheal intubation is considered the gold standard for protecting the airway.<sup>[4]</sup> However, the success rates for intubation are variable depending on airway structure, a patient's clinical status, practitioner's skills, and so forth.<sup>[5,6]</sup> The European Resuscitation Council guidelines for cardiopulmonary resuscitation (2010) recommend that

intubation be performed by experienced and trained personnel only.<sup>[7]</sup>

Endotracheal intubation is a definitive way of securing the airway and is routinely done by laryngoscopy and visualisation of cords. However, this involves distortion of upper airway to bring glottis into the line of sight<sup>[8]</sup> and in some situations such as high larynx, facial trauma, etc., tracheal intubation fails. Sufficient anatomical distortion is not always easy or possible, leading to intubation difficulties in 1-3%<sup>[9]</sup> and failure in 0.05-0.2% of cases.<sup>[10,11]</sup>

As a result of studies regarding the provision of an airway that is less invasive than intubation but safer than mask to maintain the patency of airway after anaesthesia

induction in brief surgical interventions, supraglottic airway devices (SADs) have been introduced into practice. They are inserted into glottic entry via the oral route and can be used in emergency conditions when tracheal intubation and mask anesthesia are challenging.<sup>[12]</sup>

The laryngeal mask airway and similar supraglottic airway devices use an inflatable cuff to wedge into the upper esophagus and provide a perilaryngeal seal <sup>[1]</sup>. The mask shape of these devices resembles a wedge-shaped doughnut in overall design. They have a tapered leading tip, a rounded proximal shape and with inflation these masks have a flat face when viewed from a lateral perspective. Inflatable masks provide an airway seal but can negatively impact on how these devices are inserted, how they are positioned and how they perform. On insertion the deflated leading edge of the mask can catch the epiglottis edge and cause it to down-fold or impede proper placement beneath the tongue. The best performance of the laryngeal mask airway occurs with semi-inflation <sup>[2-4]</sup>. Inflation using the recommended volume increases mask rigidity, decreases conformity with peri-laryngeal structures and lessens the effective seal pressure<sup>[2-4]</sup>. Mechanically, inflation can cause movement of the device because the distal wedge shape of the mask is forced out of the upper esophagus. Inflatable masks also have the potential to cause tissue distortion, venous compression and nerve injury <sup>[5-7]</sup>. The laryngeal mask airway and similar supraglottic airway devices use an inflatable cuff to wedge into the upper esophagus and provide a perilaryngeal seal <sup>[1]</sup>. The mask shape of these devices resembles a wedge-shaped doughnut in overall design. They have a tapered leading tip, a rounded proximal shape and with inflation these masks have a flat face when viewed from a lateral perspective. Inflatable masks provide an airway seal but can negatively impact on how these devices are inserted, how they are positioned and how they perform. On insertion the deflated leading edge of the mask can catch the epiglottis edge and cause it to down-fold or impede proper placement beneath the tongue. The best performance of the laryngeal mask airway occurs with semi-inflation <sup>[2-4]</sup>. Inflation using the recommended volume increases mask rigidity, decreases conformity with peri-laryngeal structures and lessens the effective seal pressure<sup>[2-4]</sup>. Mechanically, inflation can cause movement of the device because the distal wedge shape of the mask is forced out of the upper esophagus. Inflatable masks also have the potential to cause tissue distortion, venous compression and nerve injury <sup>[5-7]</sup>. The laryngeal mask airway and similar supraglottic airway devices use an inflatable cuff to wedge into the upper esophagus and provide a perilaryngeal seal.

SAD has the advantage of easier use without examining the vocal cords in a difficult airway. While tracheal intubation requires effort and time to maintain skills, SADs do not.<sup>[13]</sup> However, SADs are limited in their ability to completely maintain and protect the airway. For

this reason, blind intubation is required through a SAD after insertion. Many SADs have been developed and many studies have evaluated blind intubation through SADs.<sup>[14]</sup>

Tracheal intubation with Macintosh laryngoscope was considered as the “gold standard” in airway management.<sup>[15]</sup> Supraglottic airway devices (SADs) are helpful in difficult airways and in emergency life threatening situations. The use of supraglottic devices as a means of rescue in patients who are difficult to intubate or ventilate has increased in the field of anesthesiology and in emergency medicine.

Laryngeal mask airway (LMA) classic (c-LMA)<sup>[16]</sup> is one such device which is included in Difficult Airway Society guidelines for unanticipated difficult intubation.<sup>[17]</sup> Laryngeal mask airway classic was designed for maintenance of airway in emergency situations, especially by untrained personnel. Later it was modified into intubating LMA (ILMA) or LMA Fastrach.<sup>[8]</sup>

I-gel is a relatively new single-use SAD which does not have an inflatable cuff.<sup>[18-22]</sup> It is made from a soft, gel-like and transparent thermoplastic elastomer (styrene ethylene butadiene styrene) which creates a non-inflatable seal which is a mirror impression of the supraglottic anatomy.<sup>[23-25]</sup> The i-gel has several other useful design features including a gastric channel, an epiglottic ridge and a ridged flattened stem to aid insertion and reduce the risk of axial rotation.<sup>[26]</sup> The stem of the i-gel is less flexible than that of the LMA-classic and has an integral bite.<sup>[27]</sup>

I-gel results in higher sealing pressures by matching the peripharyngeal anatomy despite the absence of an inflatable cuff.<sup>[28]</sup> In a cadaveric study full glottis view was obtained in 60% of the cases soon after I-Gel insertion while some glottic opening was visible in 95% of the cases.<sup>[29]</sup> The present study was conducted at comparative evaluation of clinical performance of I-GEL with LMA fastrach in elective surgeries.

## MATERIALS AND METHOD

The present study entitled “A Prospective randomized comparison of clinical performance of I-GEL with LMA fastrach in elective surgeries” was conducted after clearance from Board of Studies, Department of Anesthesiology and Ethical committee in the Department of Anesthesiology, Teerthanker Mahaveer Medical College & Research Centre, Moradabad during the period 2015-2017, on 80 patients undergoing elective surgery under general anesthesia.

### Inclusion criteria

- Patients undergoing general anesthesia for elective surgical procedures lasting more than 30 minutes.
- American society of anesthesiology grades I
- Age group 20 to 50 years of either sex.
- BMI 18-24 Kg/m<sup>2</sup>
- Adequate mouth opening.

### Exclusion Criteria

- Patient contraindication to spinal anesthesia.
- Patients with neurological deficits, spinal cord deformities, psychological illness, hypertensive or hypovolemic.
- Emergency lower segment caesarean section.
- Patients with use of oral opioids or non-steroidal anti-inflammatory drugs.

### Sample Size

The sample size calculation was based on a 28 seconds (50%) reduction in the intubation time comparing I-gel and LMA fastrach. The sample size was calculated to be 36 patients each of I-gel and LMA fastrach to detect a significant difference between the groups at 5% alpha error and 80% power. Considering a drop-out rate of 10%, the sample size was taken to be 40 patients per group.

### Randomization

The patients were divided into two groups: Group I: Anesthesia was delivered using I-gel in 40 patients and Group LF: Anesthesia was delivered using LMA fastrach in 40 patients. Learning curve was achieved by doing 10 intubations using each of the devices on patients, prior to start of study.

### Technique of Anesthesia

Insertion of I-gel/LMA fastrach was carried out as per the study protocol. LMA fastrach was chosen according to the weight of the patient. Recommended size of LMA according to weight of patients are as follows:

#### For I-Gel/Pro-Seal Group

Size of supraglottic devices	Patients' body weight	ETT size internal diameter (mm)
I-GEL		
Size 3	30-50kg	7.0
Size 4	50-90kg	7.5
ILMA		
Size 3	30-50kg	7.0
Size 4	50-70kg	7.5
Size 5	>70kg	7.5

Lubrication of the front and back of the SAD and a jaw lift was carried out with head in neutral/extended position to facilitate its insertion. After insertion, the cuff was inflated, and its pressure adjusted to between 60-70 cmH<sub>2</sub>O. Proper placement was confirmed by listening for signs of a leak, observing rising of the chest, by auscultation and noting the presence of a normal square wave pattern on capnograph tracing, under manually assisted ventilation. Following successful insertion of the airway devices, breathing circuit was attached and patient was maintained on O<sub>2</sub>, Nitrous Oxide, Isoflurane (1%) and Intermittent doses of Inj.

Vecuronium (0.02 mg/kg IV). Surgery was allowed to commence only after the collection of the last haemodynamic data at 10 minutes post-insertion interval.

### Parameters recorded

The parameters recorded were Time taken for successful placement, Time taken for insertion of Endo-tracheal Tube, Number of insertion attempts, Quality of ventilation during anesthesia, Haemodynamic parameters such as pulse rate, systolic and diastolic blood pressure and SpO<sub>2</sub>, Airway trauma and Gastric distension.

### RESULTS

The study population consisted of 36 (45.1%) males and 44 (55.0%) females. Among I-gel group, there were 13 (32.5%) males and 27 (67.5%) females. Among LMA FASTRACH group, there were 23 (57.5%) males and 17 (42.5%) females. The mean age of the study population was 36.04±9.62 years. The mean age of the subjects in the I-gel group was 35.90±10.69 years and LMA Fastrach was 36.18±8.55.

There was no significant difference in mean Heart Rate at Baseline, 1 minute, 3 minutes and 5 minutes between I-GEL and LMA FASTRACH groups. Among I-GEL group, the mean Heart Rate was significantly more at 3 minutes in comparison to all other time intervals. Among LMA FASTRACH group, the mean Heart Rate increased significantly from baseline and 1 minute to 3 minutes which increased significantly to 5 minutes. (Table 1)

There was no significant difference in mean Systolic blood pressure at Baseline, 1 minute, 3 minutes and 5 minutes between I-GEL and LMA FASTRACH groups. Among I-GEL group, the mean Systolic blood pressure increased significantly from baseline to 1 minute and then decreased significantly to 5 minutes. Among LMA FASTRACH group, the mean Systolic blood pressure increased significantly from baseline to 1 minute and then decreased significantly to 3 and 5 minutes. (Table 2)

There was no significant difference in mean Diastolic blood pressure at Baseline, 1 minute, 3 minutes and 5 minutes between I-GEL and LMA FASTRACH groups. Among I-GEL group, there were no significant changes in the mean Diastolic blood pressure over the different time intervals. Among LMA FASTRACH group, the mean Diastolic blood pressure increased significantly from baseline to 1 minute and then decreased significantly to 5 minutes. (Table 3)

There was no significant difference in mean Arterial pressure at Baseline, 1 minute, 3 minutes and 5 minutes between I-gel and LMA FASTRACH groups. Among I-GEL group, the mean Arterial Pressure increased significantly from baseline to 1 minute and then decreased significantly to 3 and 5 minutes. Among LMA FASTRACH group, the mean Arterial Pressure increased significantly from baseline to 1 minute and then decreased significantly to 3 and 5 minutes. (Table 4)

The mean Time taken for supra-glottic device was significantly more among LMA FASTERACH (31.57±3.08) in comparison to I-gel (15.07±1.65).The mean Time taken for Endo-tracheal Tube was significantly more among I-gel (24.00±0.31) minutes in comparison to LMA FASTERACH (19.95±0.37). (Table 5)

The distribution of Number of Attempts for insertion of supra-glottic device and Number of Attempts for

Endotracheal Intubation was done between I-GEL and LMA FASTERACH groups using the Mann-whitney U-test. No significant difference was found in the distribution of number of Attempts and Number of Attempts for Endotracheal Intubation for supra-glottic device between I-GEL and LMA FASTERACH groups.(Table 6)

**Table 1: Comparison of mean Heart rate between I-gel and LMA FASTERACH groups at Baseline, 1 minute, at 3 minutes and at 5 minutes**

Heart Rate		I-GEL		LMA FASTERACH		t-test value	p-value <sup>a</sup>
		Mean	Std. Deviation	Mean	Std. Deviation		
1.	Baseline	85.70	4.44	83.70	4.77	1.941	0.056 <sup>#</sup>
2.	1 minute	86.58	4.91	83.33	4.83	2.984	0.074 <sup>#</sup>
3.	3 minutes	95.98	7.46	90.23	4.02	1.748	0.060 <sup>#</sup>
4.	5 minutes	82.53	3.73	78.73	2.77	2.784	0.058 <sup>#</sup>
p-value <sup>b</sup>		< 0.001*		< 0.001*			
Post-hoc comparisons <sup>c</sup>		3>1,2,4		3>1,2,4			

<sup>a</sup>Mann-whitney U-test  
\* Significant difference

<sup>b</sup>Friedman’s test  
<sup>#</sup> Non-significant difference

<sup>c</sup>Wilcoxon-sign rank test

**Table 2: Comparison of mean Systolic blood pressure between I-gel and LMA FASTERACH groups at Baseline, 1 minute, at 3 minutes and at 5 minutes**

Systolic blood pressure		I-GEL		LMA FASTERACH		t-test value	p-value <sup>a</sup>
		Mean	Std. Deviation	Mean	Std. Deviation		
1.	Baseline	121.25	3.67	121.53	3.97	-0.321	0.749 <sup>#</sup>
2.	1 minute	127.93	2.16	129.53	6.85	1.391	0.102 <sup>#</sup>
3.	3 minutes	120.53	2.00	122.83	4.01	2.863	0.091 <sup>#</sup>
4.	5 minutes	117.73	3.58	116.25	3.03	1.989	0.051 <sup>#</sup>
p-value <sup>b</sup>		< 0.001*		< 0.001*			
Post-hoc comparisons <sup>c</sup>		2>1,3,4		2>1,3,4			

<sup>a</sup>Mann-whitney U-test  
\* Significant difference

<sup>b</sup>Friedman’s test  
<sup>#</sup> Non-significant difference

<sup>c</sup>Wilcoxon-sign rank test

**Table 3: Comparison of mean Diastolic blood pressure between I-gel and LMA FASTERACH groups at Baseline, at 1 minute, at 3 minutes and at 5 minutes**

Diastolic blood pressure		I-GEL		LMA FASTERACH		t-test value	p-value
		Mean	Std. Deviation	Mean	Std. Deviation		
1.	Baseline	78.48	4.80	79.33	3.68	-0.889	0.377 <sup>#</sup>
2.	1 minute	81.48	2.66	84.00	2.98	-2.909	0.060 <sup>#</sup>
3.	3 minutes	80.10	4.93	81.50	3.80	-2.454	0.081 <sup>#</sup>
4.	5 minutes	75.35	3.30	76.00	4.16	-1.157	0.102 <sup>#</sup>
p-value <sup>b</sup>		0.102 <sup>#</sup>		< 0.001*			
Post-hoc comparisons <sup>c</sup>		N/A		2>1,4			

<sup>a</sup>Mann-whitney U-test  
\* Significant difference

<sup>b</sup>Friedman’s test  
<sup>#</sup> Non-significant difference

<sup>c</sup>Wilcoxon-sign rank test

**Table 4: Comparison of mean Arterial pressure between I-gel and LMA FOSTRACH groups at Baseline, at 1 minute, at 3 minutes and at 5 minutes**

Mean Arterial pressure		I-GEL		LMA FOSTRACH		t-test value	p-value
		Mean	Std. Deviation	Mean	Std. Deviation		
1.	Baseline	92.73	3.46	93.39	3.33	-0.867	0.389 <sup>#</sup>
2.	1 minute	99.29	2.09	99.18	2.20	1.327	0.083 <sup>#</sup>
3.	3 minutes	95.58	3.69	95.28	2.07	0.448	0.655 <sup>#</sup>
4.	5 minutes	88.84	3.25	89.42	2.45	-1.981	0.071 <sup>#</sup>
p-value <sup>b</sup>		< 0.001*		< 0.001*			
Post-hoc comparisons <sup>c</sup>		2>1,3,4		2>1,3,4			

<sup>a</sup>Mann-whitney U-test

<sup>b</sup>Friedman’s test

<sup>c</sup>Wilcoxon-sign rank test

\* Significant difference

<sup>#</sup> Non-significant difference

**Table 5: Comparison of mean Time taken for insertion of supra-glottic device between I-gel and LMA FOSTRACH groups**

	I-GEL		LMA FOSTRACH		t-test value	p-value
	Mean	SD	Mean	SD		
Time taken for supra-glottic device	15.07	1.65	31.57	3.08	-29.856	0.001*
Time taken for Endo-tracheal Tube	24.00	0.31	19.95	0.37	23.299	0.001*

<sup>a</sup>Mann-whitney U-test

\* Significant difference

**Table 6: Comparison of frequency distribution of Number of Attempts for supra-glottic device insertion between I-gel and LMA FOSTRACH groups**

		Groups		Total	p-value <sup>c</sup>
		I-GEL	LMA FOSTRACH		
Number of Attempts for supra-glottic device	One	40	37	77	<b>0.077<sup>#</sup></b>
		100.0%	92.5%	96.3%	
	Two	0	3	3	
		0.0%	7.5%	3.8%	
Number of Attempts for Endo tracheal Intubation	One	26	32	58	<b>0.133<sup>#</sup></b>
		65.0%	80.0%	72.5%	
	Two	14	8	22	
		35.0%	20.0%	27.5%	

<sup>c</sup>Chi-square test

<sup>#</sup> Non-significant difference

**DISCUSSION**

SGA is an integral part of difficult airway algorithm and resuscitation protocols.[<sup>22</sup>]It is also commonly used as a rescue device when a “cannot intubate, cannot ventilate” scenario arises.[<sup>23</sup>]Recently, the resuscitation outcomes such as survival to hospital discharge, return of spontaneous circulation, and 24-hour survival were shown to be better following endotracheal intubation in comparison to SGA when used for OHCA.[<sup>24,25</sup>]

In this study, overall success rate of insertion of supraglottic devices in both the groups was 100% which

was similar to various previously conducted studies. In the present study, first- attempt success rate for blind tracheal intubation was comparable in both the groups and overall success rate was higher in I group (100.0%)as compared to F group (92.5%) which was similar to the studies by Sahi et al.[<sup>26</sup>] the insertion success rate in group I-gel was 83.3% for the 1st attempt while in Group LMA Fastrach, it was 65% for the 1st attempt and Bhandari et al.[<sup>27</sup>] in which 95% reported first time and 100% overall success rate with I-gel. This was dissimilar to the results of Halwagi et

al.[<sup>28</sup>]and Sastre et al.[<sup>29</sup>]who noticed higher success rate of blind tracheal intubation with ILMA.

Raggazi et al[<sup>30</sup>]in their study found that LMA supreme has fewer insertion failures as compared to I-Gel but because of its inflatable cuff caused transient pharyngolaryngeal pain. Theiler et al.[<sup>31</sup>] in their study concluded that both LMA supreme and I-Gel have a similar insertion success and clinical performance in the simulated difficult airway situation. However, Singh et al[<sup>32</sup>]found that that I-Gel was easier to insert and required less attempts of insertion when compared with proseal LMA.

The overall intubation success rate using LMA Fastrach was comparable to published studies.[<sup>33-36</sup>]The cases in which blind tracheal intubation failed only two patients needed stylet for intubation with Macintosh laryngoscope in group I-gel and none in group LMA Fastrach. The easier and a quicker insertion of i-gel was probably due to non-inflation of cuff. Time was not wasted in inflating the cuff, and moreover, the rigid structure of LMA Fastrach causes delay in insertion as compared to i-gel.

Michalek et al.[<sup>20</sup>]did blind tracheal intubation in three different airway manikins through the i-gel with a success rate of 51%. Theiler et al.[<sup>31</sup>] studied “visualised blind intubation” through the i-gel and the LMA Fastrach. Their results showed a poor success rate (15%) with i-gel as compared with the LMA Fastrach (69%). Sastre et al.[<sup>29</sup>] also showed an inferior intubation rate of 40% through i-gel as compared to 70% with LMA Fastrach.

The mean Time taken for supra-glottic device was significantly more among LMA FASTER (31.57±3.08) in comparison to I-gel (15.07±1.65).This was similar to the studies by Moore et al.[<sup>37</sup>] the time required for tracheal intubation were significantly lower in the IG group (30 ± 11 seconds vs 50 ± 21 seconds; P < 0.0001), Chauhan *et al.*[<sup>38</sup>]have observed significantly lower insertion times with i-gel (11.12 ± 1.814 seconds) when compared with LMA proseal (15.13 ± 2.91 seconds) and Halwagi et al.,[<sup>28</sup>] showed longer intubation times with ILMA in comparison to I-Gel.

However, in the study by Sahi et al.[<sup>39</sup>]the time taken for Endotracheal intubation by Fastrach LMA was 18.953±0.925 seconds for one attempt with overall mean of 21.509±5.374 seconds while that of i-gel was 23.00±1.433 seconds for one attempt with overall mean of 26.906±7.517 seconds showing that intubation through Fastrach LMA took lesser time than i-gel, which is found to be statistically highly significant, Fernández *et al.*[<sup>40</sup>]had observed longer insertion time (32.5 seconds) with i-gel compared to LMA-S (27.1 seconds) and lower first attempt placement rates with i-gel (86%) compared to with LMA-S (95.2%) and Theiler *et al.*[<sup>31</sup>] LMA-S needed shorter insertion time (34±12 s vs. 42±23 s, P = 0.024).Theiler *et al.*[<sup>31</sup>] have attributed the longer insertion time for i-gel to the bulky design of the airway device.

The mean Time taken for Endo-tracheal Tube was significantly more among I-gel(24.00±0.31) minutes in comparison to LMA FASTER (19.95±0.37).This was similar to the study by Kapoor et al.[<sup>14</sup>](24.04 ± 9.42 seconds) and Halwagi et al(22 ± 13 seconds).[<sup>28</sup>]

The flexible silicon tipped tube is a well-designed, straight, soft, wire-reinforced silicon tube which lacks wire reinforcement in the distal inch and terminates like a conical soft tip for use with ILMA. This combined with the enhanced curved shape of the ILMA leads these flexible tubes towards the plane of the glottis at an angle of 35°. The relatively straight shape of the I-gel stem and the ending of the airway channel deep into the bowl of the cuff may direct the soft tip of FST posteriorly thereby increasing the risk of oesophageal intubation or snaring on the arytenoids. The more rigid PVC tubes have a fixed curvature directed anteriorly thereby better aligning the tube towards laryngeal inlet than FST when advanced through an I-Gel.[<sup>41</sup>]

The incidence of postoperative complications was comparable in both the groups. In the present study, dysphonia was more in I group which was similar to study conducted by Sastre et al.[<sup>29</sup>]While the incidence of sore throat was lesser in I group when compared to F group; this observation is similar to that of Keijzer *et al.*[<sup>42</sup>]

There was no significant difference in mean Heart Rate, systolic blood pressure, diastolic blood pressure and mean Arterial pressure between I-GEL and LMA FASTER groups at all time intervals. This was similar to the study by Sahi et al.[<sup>39</sup>] hemodynamic changes were comparable as is showed by insignificant statistical difference during induction, SAD insertion, intubation and throughout the surgery.

In the study by Gupta et al.[<sup>43</sup>]the mean heart rate, systolic, diastolic, mean arterial blood pressure and Spo2 in LMA-Supreme group and I-gel group did not have significant difference between two groups at different intervals of time. Shin et al.[<sup>44</sup>]found no difference in the hemodynamic data immediately after insertion of device.

Shin et al.[<sup>44</sup>]also concluded that the tongue, lip & dental trauma and blood staining of the device was more with LMA-Supreme than with I-gel but with no statistical significance. These observations are consistent with our results and with Helmy et al.[<sup>36</sup>] study in which they concluded that there was no statistically significant difference found between both I-gel and classical laryngeal mask airway groups with regard to sore throat, hoarseness and dysphonia 24 hours after the surgery.

As such, no post-operative complications were reported in the present study. The similar results were also reported by Goyal et al.[<sup>45</sup>] no sore throat and hoarseness was reported though there was blood contamination in all three SADs (i-gel, proseal, and classical). Similar to our findings, Shin et al.[<sup>44</sup>]did not determine any blood contamination or sore throat in the i-gel group who underwent orthopedic surgery in the supine position.

Uppal et al.<sup>[20]</sup> compared the i-gel with a tracheal tube, they found 12% blood contamination in relation with the insertion method and ease. Ragazzi et al.<sup>[30]</sup> compared target-controlled anesthesia with the I-gel and supreme and found one blood contamination in the I-gel group and two in the supreme group. The gel-like cuff minimizes trauma of the airway and neurovascular compression.

Theiler et al.<sup>[31]</sup> reported that Fiberoptic view of the glottis was remarkably good through the i-gel<sup>™</sup> compared with the LMA-S<sup>™</sup>. This finding and the smaller proportion of epiglottic down folding were the only statistically significant differences in favour of the i-gel<sup>™</sup>. Similar findings have been reported in the earlier fiberoptic findings<sup>[19]</sup>. Neither epiglottic down folding nor fiberoptic view could be correlated to ventilation success and possible tidal volume applied.

However, in LMA Fastrach, there was no difference in successful blind tracheal intubation with conventional tracheal tube and silicon wire reinforced tracheal tube in studies conducted by Lu et al.<sup>[11]</sup> and Kundra et al.<sup>[36]</sup> but in case of i-gel further studies are required.

## CONCLUSION

It can therefore be inferred that both Fastrach LMA and i-gel are suitable devices to be used as conduit to endotracheal intubation particularly in susceptible patients where hemodynamic disturbances during intubation are not required.

## REFERENCES

- Brain AI, Verghese C, Addy EV, Kapila A. The intubating laryngeal mask. I: Development of a new device for intubation of the trachea. *Br J Anaesth*. 1997;79:699–703.
- Lee JR, Kim MS, Kim JT, Byon HJ, Park YH, Kim HS, et al. A randomised trial comparing the i-gel (TM) with the LMA Classic (TM) in children. *Anaesthesia*. 2012;67:606–11.
- Michalek P, Hodgkinson P, Donaldson W. Fiberoptic intubation through an i-gel supraglottic airway in two patients with predicted difficult airway and intellectual disability. *AnesthAnalg*. 2008;106:1501–4.
- 7th ed. Wokingham, UK: Intersurgical Ltd; 2009. i. gel User Guide.
- Levitan RM, Kinkle WC. Initial anatomic investigations of the i-gel airway: A novel supraglottic airway without inflatable cuff. *Anaesthesia*. 2005;60:1022–6.
- Uppal V, Gangaiah S, Fletcher G, Kinsella J. Randomized crossover comparison between the i-gel and the LMA-Unique in anaesthetized, paralysed adults. *Br J Anaesth*. 2009;103:882–5.
- Brain AI. The laryngeal mask – A new concept in airway management. *Br J Anaesth*. 1983;55:801–5.
- Henderson JJ, Popat MT, Latto IP, Pearce AC. Difficult Airway Society. Difficult Airway Society guidelines for management of the unanticipated difficult intubation. *Anaesthesia*. 2004;59:675–94.
- Beylacq L, Bordes M, Semjen F, Cros AM. The I-gel, a single-use supraglottic airway device with a non-inflatable cuff and an esophageal vent: an observational study in children. *ActaAnaesthesiol Scand*. 2009;53:376–379.
- Lee JR, Kim MS, Kim JT, Byon HJ, Park YH, Kim HS, et al. A randomised trial comparing the i-gel (TM) with the LMA Classic (TM) in children. *Anaesthesia*. 2012;67:606–11.
- Theiler LG, Kleine-Brueggeney M, Luepold B, Stucki F, Seiler S, Urwyler N, et al. Performance of the pediatric-sized i-gel compared with the Ambu Aura Once laryngeal mask in anesthetized and ventilated children. *Anesthesiology*. 2011;115:102–110.
- Keller C, Brimacombe JR, Keller K, Morris R. Comparison of four methods for assessing airway sealing pressure with the laryngeal mask airway in adult patients. *Br J Anaesth*. 1999;82:286–287.
- Lopez-Gil M, Brimacombe J, Keller C. A comparison of four methods for assessing oropharyngeal leak pressure with the laryngeal mask airway (LMA) in paediatric patients. *PaediatrAnaesth*. 2001;11:319–321.
- Hughes C, Place K, Berg S, Mason D. A clinical evaluation of the I-gel<sup>™</sup> supraglottic airway device in children. *PaediatrAnaesth*. 2012;22:765–771.
- Kim MS, Lee JR, Shin YS, et al. Comparison of 2 cuff inflation methods of laryngeal mask airway Classic for safe use without cuff manometer in adults. *Am J Emerg Med* 2014;32:237–42.
- Ragazzi R, Finessi L, Farinelli I, et al. LMA Supreme vs i-gel—a comparison of insertion success in novices. *Anaesthesia* 2012;67:384–8.
- Cook TM, Green C, McGrath J, et al. Evaluation of four airway training manikins as patient simulators for the insertion of single use laryngeal mask airways. *Anaesthesia* 2007;62:713–8.
- Robak O, Leonardelli M, Zedtwitz-Liebenstein K, et al. Feasibility and speed of insertion of seven supraglottic airway devices under simulated airway conditions. *CJEM* 2012;14:330–4. [PubMed]
- Leventis C, Chalkias A, Sampanis MA, et al. Emergency airway management by paramedics: comparison between standard endotracheal intubation, laryngeal mask airway, and I-gel. *Eur J Emerg Med* 2014;21:371–3.
- Lee DW, Kang MJ, Kim YH, et al. Performance of intubation with 4 different airway devices by unskilled rescuers: manikin study. *Am J Emerg Med* 2015;33:691–6.
- Revez L, Buendia JA, Tellez D. Chemoprophylaxis in contacts of patients with leprosy: systematic review and meta-analysis. *Rev PanamSaludPublica* 2009;26:341–9.
- Kwak DI, Yoo JH, No HW, et al. Comparison between the LMA Classic<sup>™</sup> and new I-gel<sup>™</sup> supraglottic airway device: a manikin study. *Hong Kong J Emerg Med* 2013;20:25–33.
- Adelborg K, Al-Mashhadi RH, Nielsen LH, et al. A randomised crossover comparison of manikin ventilation through Soft Seal®, i-gel and AuraOnce<sup>™</sup> supraglottic airway devices by surf lifeguards. *Anaesthesia* 2014;69:343–7.
- Komasawa N, Ueki R, Kaminoh Y, et al. Evaluation of chest compression effect on airway management with air-Q®, aura-i®, i-gel®, and Fastrack® intubating supraglottic devices by novice physicians: a randomized crossover simulation study. *J Anesth* 2014;28:676–80.
- Deakin CD, Nolan JP, Soar J, et al. European Resuscitation Council guidelines for resuscitation 2010 section 4. Adult advanced life support. *Resuscitation* 2010;81:1305–52.
- Kim MS, Oh JT, Min JY, et al. A randomised comparison of the i-gel and the laryngeal mask airway classic in infants. *Anaesthesia* 2014;69:362–7.

27. Singh J, Yadav MK, Marahatta SB, Shrestha BL. Randomized crossover comparison of the laryngeal mask airway classic with i-gel laryngeal mask airway in the management of difficult airway in post burn neck contracture patients. *Indian J Anaesth.* 2012;56:348–52.
28. Radhika KS, Sripriya R, Ravishankar M, Hemanth Kumar VR, Jaya V, Parthasarathy S. Assessment of suitability of i-gel and laryngeal mask airway-supreme for controlled ventilation in anesthetized paralyzed patients: A prospective randomized trial. *Anesth Essays Res.* 2016;10:88–93. [PMC free article] [PubMed]
29. Saran S, Mishra SK, Badhe AS, Vasudevan A, Elakkumanan LB, Mishra G. Comparison of i-gel supraglottic airway and LMA-ProSeal<sup>™</sup> in pediatric patients under controlled ventilation. *J Anaesthesiol Clin Pharmacol.* 2014;30:195–8.
30. Theiler LG, Kleine-Brueggene M, Kaiser D, Urwyler N, Luyet C, Vogt A, et al. Crossover comparison of the laryngeal mask supreme and the i-gel in simulated difficult airway scenario in anesthetized patients. *Anesthesiology.* 2009;111:55–62.
31. Jindal P, Rizvi A, Sharma JP. Is I-gel a new revolution among supraglottic airway devices? – A comparative evaluation. *Middle East J Anaesthesiol.* 2009;20:53–8.
32. Atef HM, Fattah SA, Gaffer ME, Al Rahman AA. Perfusion index versus non-invasive hemodynamic parameters during insertion of i-gel, classic laryngeal mask airway and endotracheal tube. *Indian J Anaesth.* 2013;57:156–62.
33. Ali A, Ali L, Sheikh NA, Siddique SA. Airway device; Comparison of I-gel supraglottic with laryngeal mask airway for ease of insertion. *Professional Med J Dec.* 2010;17:643–7.
34. Jindal P, Rizvi AA, Khurana G, Sharma P. Safety and efficacy of insertion of supraglottic devices in anaesthetised patients by first time users. *South Afr J AnaesthesiolAnalg.* 2010;16:23–6.
35. Atef HM, Helmy AM, El-Taher EM, Henidak AM. Comparative study between I-gel, a new supraglottic airway device, and classical laryngeal mask airway in anesthetized spontaneously ventilated patients. *Middle East J Anaesthesiol.* 2012;21:583–90.
36. Helmy AM, Atef HM, El-Taher EM, Henidak AM. Comparative study between I-gel, a new supraglottic airway device and classical laryngeal mask airway in anaesthetized spontaneously ventilated patients. *Saudi J Anaesth.* 2010;4:131–6.
37. Reza Hashemian SM, Nouraei N, Razavi SS, Zaker E, Jafari A, Eftekhari P, et al. Comparison of i-gel<sup>™</sup> and laryngeal mask airway in anesthetized paralyzed patients. *Int J Crit Illn Inj Sci.* 2014;4:288–92.
38. Singh I, Gupta M, Tandon M. Comparison of clinical performance of I-Gel with LMA-ProSeal in elective surgeries. *Indian J Anaesth.* 2009;53:302–5.
39. Gabbott DA, Beringer R. The igel<sup>™</sup> supraglottic airway: A potential role for resuscitation? *Resuscitation.* 2007;73:161–2.
40. Uppal V, Gangaiah S, Fletcher G, Kinsella J. Randomized crossover comparison between the I-gel and the LMA-Unique in anaesthetized, paralysed adults. *Br J Anaesth.* 2009;103:882–5.
41. Durrani HD, Butt KJ, Sadaf S, Rehan A, Khan AM, Umar A. Comparison of LMA classic and I-gel in anesthetized, spontaneously breathing patients during elective surgical procedures. *Anaesth Pain Intensive Care.* 2013;17:274–8.
42. Siddiqui AS, Raees US, Siddiqui SZ, Haider S, Raza SA. Comparison of performance and safety of I-gel with laryngeal mask airway (classic) for general anaesthesia with controlled ventilation. *Anaesth Pain Intensive Care.* 2010;14:17–20.
43. Revi N, Harikishore, Puthur B, Ershad A comparative study on cardiovascular response and ease of insertion in classical laryngeal mask airway, ProSeal laryngeal mask airway and I-gel during surgery under general anaesthesia. *J Evid Based Med Healthc.* 2015;2:3039–46.
44. Richez B, Saltel L, Banchereau F, Torrielli R, Cros AM. A new single use supraglottic airway device with a noninflatable cuff and an esophageal vent: An observational study of the i-gel. *Anesth Analg.* 2008;106:1137–9.
45. Stewart A, Lindsay WA. Bilateral hypoglossal nerve injury following the use of the laryngeal mask airway. *Anaesthesia.* 2002;57:264–5.
46. Twigg S, Brown JM, Williams R. Swelling and cyanosis of the tongue associated with use of a laryngeal mask airway. *Anaesth Intensive Care.* 2000;28:449–50.